


My Asthma Plan

1 My usual asthma medicines

- My preventer inhaler is called _____
_____ and its colour is _____
- I take _____ puff/s of my preventer inhaler in the morning and _____ puff/s at night. I do this every day even if I feel well.
- Other asthma medicines I take every day: _____

- My reliever inhaler is called _____
_____ and its colour is _____
- I take _____ puff/s of my reliever inhaler when I wheeze or cough, my chest hurts or it's hard to breathe.
- My best peak flow is _____



If I need my blue inhaler to do any sport or activity, I need to see my doctor or asthma nurse.

2 My asthma is getting worse if...

- I wheeze or cough, my chest hurts or it's hard to breathe, **or**
- I need my reliever inhaler (usually blue) three or more times a week, **or**
- My peak flow is less than _____, **or**
- I'm waking up at night because of my asthma (this is an important sign and I will book a next day appointment)

If my asthma gets worse, I will:

- Take my preventer medicines as normal
- And also take _____ puff/s of my blue reliever inhaler every four hours
- See my doctor or nurse within 24 hours if I don't feel better



URGENT! "If your blue reliever inhaler isn't lasting for four hours you are having an asthma attack and you need to take **emergency action now (see section 3)**"



Remember to use my spacer with my inhaler if I have one.

(If I don't have one, I'll check with my doctor or nurse if it would help me)

Other things to do if my asthma is getting worse

.....

.....

.....

.....

3 I'm having an asthma attack if...

- My reliever inhaler isn't helping or I need it more than every four hours, **or**
- I can't talk, walk or eat easily, **or**
- I'm finding it hard to breathe, **or**
- I'm coughing or wheezing a lot or my chest is tight/hurts, **or**
- My peak flow is less than _____

If I have an asthma attack, I will:



Call for help



Sit up — don't lie down. Try to be calm.



Take one puff of my reliever inhaler (with my spacer if I have it) **every 30 to 60 seconds** up to a total of 10 puffs.



If I don't have my blue inhaler, or it's not helping, I need to call 999 straightaway.



While I wait for an ambulance I can use my blue reliever again, every 30 to 60 seconds (up to 10 puffs) if I need to.

Even if I start to feel better, I can't want this to happen again, so I need to see my doctor or asthma nurse today.